

Weight gain pattern of exclusively breastfed low birth weight and normal weight babies during the first six months of life.

Mohammed Faisal, Syed Manazir Ali, Maroof Jilani Khan, Azra J. Ahmed

Department of Pediatrics, J.N. Medical College, Aligarh Muslim University, Aligarh, India

Abstract

To study and compare the weight gain pattern of Exclusively Breastfed Low birth weight and normal weight babies during the first six months of life in Neonatology Section of the Department of Pediatrics of Jawaharlal Nehru Medical College, Aligarh Muslim University, Aligarh. A prospective Study was conducted between June 2004 to December 2005. Three hundred and sixty three low birth weight babies (cases) and 121 normal birth weight babies (controls) formed the basis of this study. Follow up of these babies was done during regular visits to well baby clinic over the next six months at monthly interval. All low birth weight babies born of singleton pregnancy in JN Medical college hospital without any gross congenital malformations during a period of six months and their mothers were included in the study. Controls were term appropriate for gestational age (AGA) babies born of singleton pregnancy without any complications.

The mean birth weight was 2.04 ± 0.36 kg in study group whereas it was 2.84 ± 0.29 kg in controls. The mean weight of exclusively breastfed babies in the study group was observed to be constantly below the 5th percentile at birth and also during the 6 months follow up. However in exclusively breastfed controls, the birth weight was at 10th percentile. On follow up, the weight was between 10th to 25th percentile in the first 3 months of life and then almost reached 25th percentile of NCHS in the next 3 months. An average daily weight gain of 19.83 grams and 22.79 grams was observed in study and control group respectively during the six months follow up. Exclusive breastfeeding is important to sustain better growth during first 6 months of life.

Key words: Low birth weight (LBW), Appropriate for gestational age (AGA), NCHS Centiles, Exclusive breast feeding.

Accepted June 17 2009

Introduction

The birth weight of infant is the single most important determinant of its chances of survival, healthy growth and development [1]. Low birth weight has been defined as a birth weight of less than 2.5 kg, the measurement being taken preferably within the first hour of life, before significant postnatal weight loss has occurred [2]. Low birth weight is one of the most serious challenges in maternal and child care in both developed and developing countries. Babies born with a low birth weight present unique problems pertaining to their survival and subsequent growth and development. In addition to the high risk of mortality and morbidity, these babies show a pattern of growth different from normal term babies [3]. Growth of these babies, especially head growth

has bearing on the neurodevelopmental outcome. Very low birth weight babies are at increased risk of poor long term growth and cognitive and language deficits. [4]. In India 30-40% of live births is constituted by low birth weight babies, 60-70% being term intrauterine growth retarded and the rest preterm [5]. Further low birth weight is a prospective marker of mothers nutritional and health status [6]. The objective of our study was to study and compare the weight gain pattern of Exclusively Breastfed Low birth weight and normal babies during the first six months of life.

Methods

A prospective Study was conducted between June 2004 to December 2005. Three hundred and sixty three low




Figure 1: Comparison of weight in study and control group with NCHS (50th centile)

The growth pattern in relation to mean weight and comparison with NCHS norms in both study and control group is presented in table 2. The mean birth weight was 2.04±0.36kg in study group whereas it was 2.84±0.29kg in controls. The mean weight of exclusively breastfed babies in the study group was observed to be constantly below the 5th percentile at birth and also during the 6 months follow up. However in exclusively breastfed controls, the birth weight was at 10th percentile. On follow up, the weight was between 10th to 25th percentile in the first 3 months of life and then almost reached 25th percentile of NCHS in the next 3 months. An average daily weight gain of 19.83 grams and 22.79 grams was observed in study and control group respectively during the six months follow up.

Discussion

An analysis of the breastfeeding pattern at 3 months of age reveals that 86% and 85.7% of babies were exclusively breastfed in study group and controls respectively, whereas partial breastfeeding was given only in 5.9%and 4.8% respectively. The number of exclusively breastfed infants declined to 52.9% and 78.6% at 6 months in study group and controls respectively. In 1996, study by Abdurrahiman, Ahmad [7] on effects of maternal nutritional status on the outcome of pregnancy , showed that exclusive breastfeeding was given in 80% of babies during first 4 months which declined to only 10% at six months of age. The increased incidence of exclusive breastfeeding in our study is a reflection of our efforts towards implementing a hospital breastfeeding policy.

The mean weight of babies in the study group (2.04±.36kg) was observed to be constantly below the 5th

percentile at birth and during the 6 months follow up. The pattern of serial weight gain in this group is in close agreement with serial follow up of a slum population from Mysore city observed by Manjrekar et al [8]. However in exclusively breastfed controls, the birth weight was at 10th percentile. On follow up, the weight was between 10th to 25th percentile in the first 3 months of life and then almost reached 25th percentile of NCHS in the next 3 months. These findings were similar to a 30th percentile range seen by Ghosh et al [9] in babies born to mothers belonging to upper socioeconomic group and 25th centile and below observed by Mathur et al [10] in his study on growth pattern in breastfed babies during first 6 months of life which are comparable to our figures in control group. This finding is also supported by the study by otaige BE etal [11] who showed that exclusively breastfed infants have growth patterns comparable with the NCHS standards and that the growth in weight of exclusively breastfed infants appears adequate for the first 6 months of life.

A better weight gain of approximately 630 grams per month and 710 grams per month was seen in study and control group in the first 3 months. The weight gain decreased in both study and controls after 3 months, being about 560g per month and 650g per month respectively in next 3 months. Thus growth faltering was observed in both study group and controls after 3 months of age, though more pronounced in the study group. The initial good weight gain in both study group and controls could be accounted for the high percentage of exclusive breast feeding during the first 3 months when even mothers with fairly poor nutrition secrete fairly adequate amount of milk as already has been observed by Ghai and Sandhu [12]. An average daily weight gain of 19.83 grams and 22.79 grams was observed in study and control group respectively during the six months follow up. This average daily weight gain of 19.83 grams in our study in low birth weight group is comparable to the mean velocity of weight gain of 19.78g/day for all neonates for the first 4 months in a study conducted by Mathur, Seth, Misra [13].

References

1. Chhabra S, Sharma S. Variables affecting foetal weight. Indian Journal of Maternal and Child Health. 1994; 5 (3): 61-64
2. World Health Organization. International Statistical Classification of Diseases and related health Problems, Tenth Revision. Volume 2: Instruction Manual, Geneva, WHO. 1993.
3. Babson GS. Growth of Low birth weight infants. J Pediatr 1970; 77: 11-18.
4. Ericson A, Anders; Kallen B, Bengt. Very low birth weight boys at the age of 19. Archives of Disease in

- Childhood Fetal & Neonatal Edition. 1998 May; 78: F171-174.
5. World Health Organization. Global Health situation and projections. Estimates; 1992.
 6. Sachdev HPS. Low birth weight in South Asia, Malnutrition in South Asia: A Regional Profile, Ed. Sturat Gillespie, Rosa Publication No.5, UNICEF, 1997: 23-50
 7. AbdurRahiman E, Ahmad A.J. Effect of maternal nutritional status on the outcome of pregnancy and growth during first Six months of life. Thesis submitted for MD, Department of Pediatrics, JNMCH, AMU, 1996.
 8. Manjrekar C, Vishalakshi MP, Begum NJA, Padma ON. Breast feeding ability of under nourished mother and physical development of their infants during 0-1 year. *Indian Pediatr* 1985; 22: 801-808
 9. Ghosh S, Hooja V, Ahmed H. A longitudinal study of length, weight and head circumference from birth to 2 years among children of higher socio-economic urban community in Delhi. *Indian Pediatr* 1974; 11: 395-398.
 10. Mathur S, Mathur G P, Gupta U, Singh YD, Kushwaha K P, Verma A et al .Growth pattern in breast feed babies during first six months of life. *Indian Pediatr* 1994 March; 31: 275-278.
 11. Otaigbe BE, Alikor EA, Nkanginieme KE. Growth pattern of exclusively breastfed infants in the first six months of life: A study of babies delivered at the University of Port Harcourt Teaching Hospital, Rivers State, Nigeria. *Niger J Med*, 2005; 14 (2):137-145.
 12. Ghai OP, Sandhu RK. Study of physical growth of *Indian* children in. Delhi. *Indian J Pediatr* 1968; 35; 91-108.
 13. Mathur NB, Seth A, Mishra TK .Physical growth and biochemical indicators of protein adequacy in exclusively breastfed neonates weighing less than 1500 g. *J Trop Pediatr*. 2000; 46 (5): 288-292

Correspondence:

Syed Manazir Ali
 Department of Pediatrics
 J.N. Medical College
 Aligarh Muslim University
 Aligarh 202 002 India

e-mail: manazir1958@yahoo.com